

Duck Bag

Keeping soiled instruments moist

- Instrument humidity pack
- For transport and storage prior to instrument reprocessing
- Pre moistened ready for immediate use
- 4-Zyme sachet included works at room temperature
- Heat sealed bag is easy to use & single use
- Available in 3 sizes:
 - Half DIN 420mm (16.5") wide x 470mm (18.5") deep
 - Full DIN 400mm (15.7") wide x 780mm (30.7") deep
 - Super Size 600mm (23.6") wide x 1000mm (39.3") deep



Medisafe







The pre moistened Medisafe Duck Bag[™] prevents soiled instruments from drying during transportation from point of use to CSSD. Place your soiled instruments and tray inside the see-through Duck Bag and seal to maintain a humid environment.

National/International guidelines and instrument manufacturers recommend that the reprocessing of soiled instruments begin within 30 minutes of end of use to ensure that soil, especially blood, does not dry on the surface of used instruments.

The CFPP highlights the importance of keeping instruments moist between use and reprocessing:

"Prions are hydrophobic proteins. The attachment of hydrophobic proteins to surfaces becomes less reversible if they are allowed to dry fully onto a surface. Keeping the environment around soiled instruments at or near saturation humidities (moist) prevents full attachment of hydrophobic proteins such that they are more efficiently removed by cleaning." DOH, CFPP 01-01, Page 20, March 2013



Duck Bag

Keeping soiled instruments moist




Name:	Duck Bag
Classification:	CE marked Class 1
Ingredients	pH neutral aqueous solution of protease HT and protease LT, lipase and amylase enzymes with corrosion inhibitor and surfactants
Storage condition	between 2°C and 38°C(35.6°F - 100.40°F)
рН	7.5-8.2
Shelf life	Up to 2 years - applies only to detergent
Packaging:	50 bags per box

Instruction for use Medisafe Duck Bag™



1.Empty contents of 4-Zyme sachet into Duck Bag™



2. Place instruments and tray into Duck Bag™



3. Close seal to ensure humid environment during transport and storage



4. Transport Duck Bag[™] from OR to CS



5. Remove instruments before reprocessing



6. Discard used Duck Bag™ and reprocess instruments according to manufacturer's instructions for use